



UPPSALA  
UNIVERSITET

**Vitaly Makaganiuk**

PhD student, SNSB financed

**Andreas Korn**

coordinator of the Gaia DPAC  
work package  
“Synthetic Stellar Spectra”

**Bengt Edvardsson**

representing the MARCS team

Department of Physics and  
Astronomy

Uppsala University,  
75120 Uppsala, Sweden



# Taking MARCS to the next level

## Ca IR triplet lines in NLTE

### Context


- Using the MARCS code (Gustafsson *et al.* 2008), Uppsala is producing model atmospheres for F, G and K stars ( $4000 \text{ K} \leq T_{\text{eff}} \leq 8000 \text{ K}$ ) spanning five orders of magnitude in metallicity and surface gravity.
- These model atmospheres are used to produce synthetic observables for the stellar classification to be performed in the framework of the **ESA Gaia space mission**:
  1. fluxes for optical spectro-photometry (so-called **BP/RP spectra**)
  2. spectra for near-IR spectroscopy around the Ca IR triplet lines at  $R = 11500$  (so-called **RVS spectra**)
- While the basic assumptions of, e.g., mixing-length convection and LTE cannot be relaxed globally on the timescale of the Gaia mission, we are making an effort to **remove some of the most obvious biases to sharpen Gaia’s view of the stellar content of the Galaxy**.
- Here, we report on work in progress to relax the assumption of LTE in the formation of the Ca IR triplet lines (see Fig. 1) for the whole grid of MARCS spectra.

### NLTE calculations


- The NLTE calculations are performed using an ALI version of DETAIL tailored for cool stars (Gehren, priv. comm.). The Ca I/II model atom is the one recently presented by Mashonkina, Korn & Przybilla (2007). It was carefully calibrated on stars with well-known stellar parameters and removes systematic differences between weak and strong lines of Ca I and II.

An example:

The metal-poor turnoff star **HD 84937** ( $T_{\text{eff}} = 6350 \text{ K}$ ,  $\log g = 4.0$ ,  $[\text{Fe}/\text{H}] = -2.2$ ) has  $[\text{Ca}/\text{Fe}]_{\text{LTE}} \approx 0.3$  as measured from optical Ca I lines. However, Ca II 8498 indicates  $[\text{Ca}/\text{Fe}]_{\text{LTE}} \approx 0.65$  (see Fig. 2). **Using our NLTE approach, a single  $[\text{Ca}/\text{Fe}]$  of  $0.40 \pm 0.05$  is derived**, a typical value for  $[\alpha/\text{Fe}]$  in halo stars.



**Figure 1:** Spectral range covered by Gaia’s RVS spectrograph: Sun observed with an FTS spectrograph (Kitt Peak atlas, Kurucz *et al.* 1984) and compared to the latest model spectrum produced with MARCS and BSYN.



**Figure 2:** Ca II 8498.1 in HD 84937. A good fit is obtained at the calcium abundance derived from weaker optical lines of Ca I while the LTE profile (grey) is clearly too weak. The non-LTE effect is  $-0.26$  dex in log (abundance).

### Spin-Offs

- The photospheric NLTE line strengths of the Ca IR triplet lines will serve as a point of reference in Gaia’s classification of (chromospherically) active stars (in collaboration with A. Lanzafame *et al.*, Catania, Italy)
- Optical Ca I NLTE corrections can be computed with little overhead. We plan to make these available with a web interface.
- The near-IR model spectra ( $R = 200000$ ) produced in this way may prove useful in modelling the integrated IR light of old stellar populations (e.g dSphs).