



# Cosmology AS7009, 2008 Lecture 4



## Outline

- Properties and fate of single-component Universes
    - Empty Universe
    - Flat Universe
    - Matter/Radiation/Lambda-dominated Universe
    - Einstein-de Sitter Universe
  - Properties and fate of multiple-component Universes
    - Matter + curvature
    - Benchmark model
    - Other dark energy scenarios
- Covers chapters 5 & 6 in Ryden

## What the dynamics of the Universe can tell you



## Density evolution (general)

- Friedmann equation + fluid equation + equation of state  $\rightarrow a(t), \epsilon(t)$  or  $t(a), \epsilon(a)$
- Problem: There are many components in the Universe  $\rightarrow$ 
  - Evolution complicated
  - Different components dominate evolution at different times

## Multiple energy components

Often, one just writes  $\epsilon$  and  $\Omega$

$$\epsilon_{\text{tot}} = \sum_w \epsilon_w$$

$$\Omega_{\text{tot}} = \sum_w \frac{\epsilon_w}{\epsilon_c}$$

- Remember:
- $w = 0$  (non-relativistic matter)
  - $w = 1/3$  (radiation and relativistic matter)
  - $w = -1$  (cosmological constant)
  - $w < -1/3$  (dark energy)  $\rightarrow$  Positive acceleration

## Density evolution

Fluid equation + eq. of state (see exercise session)  $\rightarrow$

$$\epsilon_w(a) = \epsilon_{w,0} a^{-3(1+w)}$$

$$\begin{cases} w_m = 0 & \Rightarrow \epsilon_m(a) = \epsilon_{m,0} a^{-3} \\ w_r = 1/3 & \Rightarrow \epsilon_r(a) = \epsilon_{r,0} a^{-4} \\ w_\Lambda = -1 & \Rightarrow \epsilon_\Lambda(a) = \epsilon_{\Lambda,0} \end{cases}$$

## The benchmark model

The currently favoured cosmological model:

$$\Omega_M \approx 0.3$$

$$\left. \begin{aligned} \Omega_M &= \Omega_{\text{Non-baryons (CDM)}} + \Omega_{\text{Baryons}} \\ \Omega_{\text{Non-baryons (CDM)}} &\approx 0.26 \\ \Omega_{\text{Baryons}} &\approx 0.04 \end{aligned} \right\} \text{More on this in the dark matter lecture}$$

$$\Omega_\Lambda \approx 0.7$$

$$\Omega_R \approx 8.4 \times 10^{-5}$$

$$\Omega_R = \Omega_{\text{CMB}} + \Omega_\nu + \Omega_{\text{starlight}}$$

$$\Omega_{\text{CMB}} \approx 5.0 \times 10^{-5}$$

$$\Omega_\nu \approx 3.4 \times 10^{-5}$$

$$\Omega_{\text{starlight}} \approx 1.5 \times 10^{-6}$$

$$\Omega_{\text{tot}} = \Omega_M + \Omega_\Lambda + \Omega_R \approx 1.0$$

→ Flat Universe ( $\kappa = 0$ )

## Why are single-component Universes relevant then?

$$w_m = 0 \Rightarrow \epsilon_m(a) = \epsilon_{m,0} a^{-3}$$

$$w_r = 1/3 \Rightarrow \epsilon_r(a) = \epsilon_{r,0} a^{-4}$$

$$w_\Lambda = -1 \Rightarrow \epsilon_\Lambda(a) = \epsilon_{\Lambda,0}$$


Values of  $\Omega_M, \Omega_\Lambda, \Omega_R$  today + these relations →

$\epsilon_M \geq \epsilon_\Lambda$  at some point in the past  
(matter-dominated Universe)

$\epsilon_R \geq \epsilon_M$  even further back  
(radiation-dominated Universe)

The different epochs of the Universe can be approximated by single-component evolution

## Different components dominate the sum at different times



## The Milne Universe (empty)

Of pretty limited relevance for our Universe, but provides simple demonstration of how to derive current age of Universe,  $a(t)$  and  $t(z)$  from the Friedmann equation

$$H(t)^2 = \frac{8\pi G}{3c^2} \epsilon(t) - \frac{\kappa c^2}{R_0^2} \frac{1}{a(t)^2}$$

$$\epsilon(t) = 0 \Rightarrow$$

$$H(t)^2 = -\frac{\kappa c^2}{R_0^2} \frac{1}{a(t)^2} \Rightarrow$$

$$\dot{a}(t)^2 = -\frac{\kappa c^2}{R_0^2}$$

## The Milne Universe (empty) II

Empty, static Universe

Empty, negatively curved

$$\dot{a}(t)^2 = -\frac{\kappa c^2}{R_0^2}$$

$$\kappa = 0 \Rightarrow \dot{a}(t) = 0$$

$$\kappa = -1 \Rightarrow$$

$$\dot{a}(t) = \frac{da}{dt} = \pm \frac{c}{R_0}$$

If expanding (not contracting)  $\Rightarrow$

$$t_0 = \frac{R_0}{c}, \quad a(t) = \frac{t}{t_0}, \quad t(z) = \frac{t_0}{1+z}$$

Current age of Universe:

## Fate of the Universe in a Milne Universe



Eternal, constant expansion ('Big Chill')

## Proper distance in a Milne Universe

Proper distance from us to object which emitted light at  $t_e$  :

$$d_p(t_0) = c \int_{t_e}^{t_0} \frac{dt}{a(t)}$$

$$a(t) = \frac{t}{t_0} \Rightarrow d_p(t_0) = ct_0 \int_{t_e}^{t_0} \frac{dt}{t} = ct_0 \ln\left(\frac{t_0}{t_e}\right)$$

$$t_e = \frac{t_0}{1+z} \Rightarrow d_p(t_0) = ct_0 \ln(1+z) = \frac{c}{H_0} \ln(1+z)$$

## Flat, single-component Universes

For  $w \neq -1$ :

$$a(t) = \left(\frac{t}{t_0}\right)^{2/(3+3w)}$$

$$\mathcal{E}(t) = \mathcal{E}_0 \left(\frac{t}{t_0}\right)^{-2}$$

## Important types of flat, single-component Universes

Radiation-only Universe (radiation-dominated epoch)

$$t_0 = \frac{1}{2H_0}, \quad a(t) = \left(\frac{t}{t_0}\right)^{1/2}$$

Matter-only Universe (matter-dominated epoch)

$$t_0 = \frac{2}{3H_0}, \quad a(t) = \left(\frac{t}{t_0}\right)^{2/3}$$

## $\Lambda$ -only Universe I

$\Lambda$  has  $w = -1 \Rightarrow$

$$\dot{a}^2 = \frac{8\pi G \mathcal{E}_\Lambda}{3c^2} a^2$$

Rearrange:

$$\dot{a}^2 = H_0^2 a^2 \quad \text{if}$$



$$H_0 = \left(\frac{8\pi G \mathcal{E}_\Lambda}{3c^2}\right)^{1/2}$$

## $\Lambda$ -only Universe II

Solution :

$$a(t) = e^{H_0(t-t_0)} \quad \text{de Sitter Universe (de Sitter phase)}$$

Same growth as in Steady state cosmology



## Einstein-de Sitter Universe I


This served as the benchmark model up until the mid-1990s

- Flat (i.e. critical-density), matter-dominated Universe


- $\Omega_M = 1.0, \Omega_{tot} = 1.0, \kappa = 0$

$$t_0 = \frac{2}{3H_0}, \quad a(t) = \left(\frac{t}{t_0}\right)^{2/3}$$

## Einstein-de Sitter Universe II




Asymptotically approaches zero expansion, but no recollapse



Einstein-de Sitter Universe

## Multiple-component Universes I

COMPOSITION OF THE COSMOS



Note: Valid for current time only!

## Multiple-component Universes II

$$H(t)^2 = \frac{8\pi G}{3c^2} \epsilon(t) - \frac{\kappa c^2}{R_0^2} \frac{1}{a(t)^2}$$

Recall:

$$\epsilon(t) = \epsilon_M(t) + \epsilon_R(t) + \epsilon_\Lambda(t) + \dots$$

## Multiple-component Universes III

For 3 components with arbitrary curvature, the FE may be rewritten:

$$\frac{H(t)^2}{H_0^2} = \underbrace{\frac{\Omega_{R,0}}{a(t)^4} + \frac{\Omega_{M,0}}{a(t)^3} + \Omega_{\Lambda,0}}_{\text{The 3 components}} + \underbrace{\frac{1 - \Omega_{tot,0}}{a(t)^2}}_{\text{Curvature}}$$

The 3 components Curvature

More components can easily be added as additional terms, as long as you know their  $a(t)$ -dependence

### Multiple-component Universes IV

Time (from Big Bang):

$$t(a) = \frac{1}{H_0} \int_0^a \frac{da}{[\Omega_{R,0}a^{-2} + \Omega_{M,0}a^{-1} + \Omega_{\Lambda,0}a + (1 - \Omega_{tot,0})]^{1/2}}$$

Lookback-time =  $t_0 - t(a)$


Nasty integral! No analytical solution in general case – must be integrated numerically!

### Special case: Matter + Curvature I

$$\frac{H(t)^2}{H_0^2} = \frac{\Omega_{M,0}}{a(t)^3} + \frac{1 - \Omega_{M,0}}{a(t)^2}$$

H(t)-evolution → Possible fates of Universe  
This is one of the hand-in exercises!  
(Note: Lots of help on page 85 in Ryden)


### Special case: Matter + Curvature II



$K = +1$        $K = 0$        $K = -1$   
'Big Crunch' Finite      'Big Chill' Infinite\*      'Big Chill' Infinite\*


\* Note: In the case of a simple topology

### Special case: Matter + Curvature III




Up until the mid-90s, these were the only 3 possible fates of the Universe typically quoted in textbooks

### Matter + curvature + $\Lambda$



### Properties of the benchmark model

- Matter-radiation equality:
    - $a \approx 2.8 \times 10^{-4}$
    - $t \approx 4.7 \times 10^4$  yr
  - Matter- $\Lambda$  equality:
    - $a \approx 0.75$
    - $t \approx 9.8$  Gyr
  - Now:
    - $a = 1$
    - $t \approx 13.7$  Gyr
- } Hand-in exercise!



### Other Dark Energy Models

What if dark energy is something other than  $\Lambda$ ?

- Varying equation of state  $w(t)$
- Constant  $w \neq -1$ ?

Future fates of the dark-energy universe

The diagram illustrates the evolution of the universe from the Big Bang to the present 'Current universe'. Three potential future paths are shown: 
 

- Big Crunch:** A path where dark energy reverses, leading to a collapse.
- Indefinite expansion:** A path where the cosmological constant remains constant, leading to continued expansion.
- Big Rip:** A path where dark energy destabilizes, leading to the tearing apart of the universe.

More on this in exercise session

Our ignorance about the nature of dark energy  $\rightarrow$   
Ultimate fate of our Universe unclear