


# Cosmology AS7009, 2011 Lecture 2




## Outline

- The cosmological principle:
  - Isotropy
  - Homogeneity
- Big Bang vs. Steady State cosmology
- Redshift and Hubble's law
- Scale factor, Hubble time, Horizon distance
- Olbers' paradox: Why is the sky dark at night?
- Particles and forces
- Theories of gravity: Einstein vs. Newton
- Cosmic curvature

Covers chapter 2 + half of chapter 3 in Ryden



## The Cosmological Principle I

- Modern cosmology is based on the assumption that the Universe is:
    - Homogeneous
    - Isotropic
- } The cosmological principle



## The Cosmological Principle II

- These tenets *seem* to hold on large scales (>100 Mpc), but definitely not on small




## The Perfect Cosmological Principle

- In this case, one assumes that the Universe on large scales is:
  - Homogeneous
  - Isotropic
  - Non-evolving

This is incompatible with the Big Bang scenario, but the *Steady State model* (popular in the 1940-1960s) was based on this idea

## Steady State Cosmology

Universe continuously expands, but due to continuous creation of matter, no dilution occurs → Steady State, no hot initial Big Bang and no initial singularity...




... But this cosmology fails to explain the CMBR,  $T_{\text{CMBR}}(z)$ , the production of light elements and the redshift evolution of galaxies & AGN

## Suggestion for Literature Exercise: Quasi-Steady State Cosmology


- Attempt in 1990s to resurrect Steady-State and explain the:
  - CMBR
  - production of light elements
  - dark matter
  - supernova type Ia data
  - Large-scale structure

## Suggestion for Literature Exercise: Quasi-Steady State Cosmology

- Cyclic creation events → Long-term steady expansion, but with short-term oscillations
- To meet observational constraints, QSS requires:
  - Strange intergalactic dust
  - Cyclic creation events ("little bangs") which cause local expansion of space




## Redshift



Definition of redshift:

$$z = \frac{\lambda_{\text{obs}} - \lambda_{\text{em}}}{\lambda_{\text{em}}}$$

## Hubble's law I



## Hubble's law II

Hubble's law:

The Hubble "constant" →  $H_0$  → Luminosity distance


$$z = \frac{H_0 d}{c}$$

In observational astronomy, the term recession velocity,  $v$ , occurs frequently:


At low  $z$ :

$$z \approx \frac{v}{c} \rightarrow v = H_0 d$$

## Expansion of the Universe I




## Expansion of the Universe II




## Redshift and distance I

- Low redshift ( $z \approx 0$ ) corresponds to:
  - Small distance (local Universe)
  - Present epoch in the history of the Universe
- High redshift corresponds to:
  - Large distance
  - Earlier epoch in the history of the Universe




## Redshift and distance II


- But beware:
  - At low redshift, Doppler components coming from peculiar motions may be substantial – must be corrected for before  $d$  is derived from  $z$  or  $v$
  - The redshift coming from cosmic expansion is not a Doppler shift – don't treat it like one!
  - The linear version of Hubble's law is only appropriate at  $z < 0.15$  (at 10% accuracy)



## Scale factor



## Scale factor and redshift



## The Hubble “constant”

$$H_0 \approx 72 \pm 5 \text{ km s}^{-1} \text{ Mpc}^{-1} [\text{s}^{-1}]$$

Today

Errorbars possibly underestimated...

Note: Sloppy astronomers often write km/s/Mpc...

In general:

$$H \equiv \frac{\dot{a}}{a}$$

Not a constant in our Universe!


## Hubble time

In the case of constant expansion rate, the Hubble time gives the age of the Universe:

$$t_H = \frac{1}{H_0} \approx 14 \text{ Gyr}$$

In more realistic scenarios, the expansion rate changes over time, but the currently favoured age of the Universe is still pretty close – around 13–14 Gyr.

## Olbers' paradox I




*"Why is the sky dark at night?"*  
(Heinrich Olbers 1926)

If the Universe is:


- Spatially infinite (i.e. infinite volume)
- Infinitely old and unevolving

- then the night sky should be bright!

## Olbers' paradox II




## Olbers' paradox III



Planet Earth surrounded by stars in an infinite, unevolving Universe

## Olbers' paradox IV



Main solution: The Universe has finite age  
The light from most stars have not had time to reach us!

## Horizon distance

- Horizon distance = Current distance to the most faraway region from which light has had time to reach us
- This delimits the causally connected part of the Universe an observer can see at any given time
- Horizon distance at time  $t_1$ :

$$d_{\text{hor}}(t_1) = c \int_{t=0}^{t_1} \frac{dt}{a(t)}$$

- Most realistic scenarios give:  
 $d_{\text{hor}}(t_0) \sim c/H_0$  (the so-called Hubble radius)

## Particles and forces I

- The particles that make up the matter we encounter in everyday life:
  - Protons, p  
938.3 MeV
  - Neutrons, n  
939.5 MeV
  - Electrons,  $e^-$   
0.511 MeV

Baryons (made of 3 quarks)

Lepton

Since most of the mass of 'ordinary matter' is contributed by protons and neutrons, such matter is often referred to as *baryonic*. Examples of mostly baryonic objects: Planets, stars, gas clouds (but not galaxies or galaxy clusters)

## Particles and forces II

- Other important particles (for this course):

- Photon,  $\gamma$   
Massless, velocity:  $c$

- Neutrinos,  $\nu_e, \nu_\mu, \nu_\tau$   
eV (?), velocity close to  $c$   
Interacts via weak nuclear force only

Leptons

## Particles and forces III

- The four forces of Nature:
  - Strong force
    - Very strong, but has short range ( $\sim 10^{-15}$  m)
    - Holds atomic nuclei together
  - Weak force
    - Weak and has short range
    - Responsible for radioactive decay and neutrino interactions
  - Electromagnetic force
    - Weak but long-range
    - Acts on matter carrying electric charge
  - Gravity
    - Weak, very long-range and always attractive

On the large scales involved in cosmology, gravity is by far the dominant one

## Newtonian gravity

- Space is Euclidian (i.e. flat)
- Planets are kept in their orbits because of the gravitational force:

$$F = -\frac{GM_g m_g}{r^2}$$

Gravitational mass

- The acceleration resulting from the gravitational force:

$$F = m_i a$$

Inertial mass

## Equivalence Principle


- Gravitational acceleration towards an object with mass  $M_g$  is:

$$a = -\frac{GM_g}{r^2} \left( \frac{m_g}{m_i} \right)$$


- Empirically  $M_g = M_i$  (to very high precision)
- The equality of gravitational mass and inertial mass is called the equivalence principle
- In Newtonian gravity,  $M_g = M_i$  is just a strange coincidence, but in General Relativity, this stems from the idea that masses cause curvature of space

## General Relativity

- 4D space-time
- Mass/energy curves space-time
- Gravity = curvature
- Pocket summary:
  - Mass/energy tells space-time how to curve
  - Curved space-time tells mass/energy how to move




## Small-scale curvature



## Global Curvature I


In the world models of general relativity, our Universe may have spatial curvature (on global scales)



## Global Curvature II


Very tricky stuff...

- This an *intrinsic* curvature in 3D space
- Note: No need for encapsulating our 3D space in 4D space to make this work



## Global Curvature III

Angles in curved spaces



## Metrics I

- Metric: A description of the distance between two points
- Metric in 2 dimensional, flat space:

$$ds^2 = dx^2 + dy^2 \quad (\text{Pythagoras})$$

- Metric in 3 dimensional, flat space:

$$ds^2 = dx^2 + dy^2 + dz^2$$

## Metrics II

- Metric in 3 dimensions, flat space, polar coordinates:

$$ds^2 = dr^2 + r^2 d\Omega^2$$

$$d\Omega^2 = d\theta^2 + \sin^2 \theta d\phi^2$$

- Metric in 3 dimensions, arbitrary curvature:

$$ds^2 = \frac{dx^2}{1 - \kappa x^2 / R^2} + x^2 d\Omega^2$$

Flat :  $\kappa = 0, x = r$

Negative :  $\kappa = -1, x = R \sinh(r/R)$

Positive :  $\kappa = 1, x = R \sin(r/R)$

Curvature radius