

Cosmology 1FA209, 2016

Lecture 3: Robertson-Walker metric & cosmic dynamics

$$\frac{\partial}{\partial \theta} M T(\xi) = \frac{\partial}{\partial \theta} \int_{\mathbb{R}_n} T(x) f(x, \theta) dx = \int_{\mathbb{R}_n} \frac{\partial}{\partial \theta} T(x) f(x, \theta) dx$$

$$\frac{\partial}{\partial a} \ln f_{a, \sigma^2}(\xi_1) = \frac{(\xi_1 - a)}{\sigma^2} f_{a, \sigma^2}(\xi_1) = \frac{1}{\sqrt{2\pi\sigma}} \exp\left\{-\frac{(\xi_1 - a)^2}{2\sigma^2}\right\} \frac{(\xi_1 - a)}{\sigma^2}$$

$$\int_{\mathbb{R}_n} T(x) \cdot \frac{\partial}{\partial \theta} f(x, \theta) dx = M\left(T(\xi) \cdot \frac{\partial}{\partial \theta} \ln L(\xi, \theta)\right) \cdot \int_{\mathbb{R}_n} \frac{\partial}{\partial \theta} T(x) f(x, \theta) dx$$

$$\int_{\mathbb{R}_n} T(x) \cdot \left(\frac{\partial}{\partial \theta} \ln L(x, \theta)\right) \cdot f(x, \theta) dx = \int_{\mathbb{R}_n} T(x) \cdot \left(\frac{\frac{\partial}{\partial \theta} f(x, \theta)}{f(x, \theta)}\right) f(x, \theta) dx$$

$$\frac{\partial}{\partial \theta} M T(\xi) = \frac{\partial}{\partial \theta} \int_{\mathbb{R}_n} T(x) f(x, \theta) dx = \int_{\mathbb{R}_n} \frac{\partial}{\partial \theta} T(x) f(x, \theta) dx$$

$$1 \cdot \exp\left\{-\frac{(\xi_1 - a)^2}{2\sigma^2}\right\} \cdot \frac{\partial}{\partial a} \ln f_{a, \sigma^2}(\xi_1) = \frac{(\xi_1 - a)}{\sigma^2}$$

Outline

- The Robertson-Walker metric
- Proper distance
- Is the Universe Infinite?
- Computational tools:
 - Friedmann equation
 - Fluid equation
 - Acceleration equation
 - Equation of state
- Cosmic dynamics
- The cosmological constant

Covers half of chapter 3 + chapter 4 in Ryden

Recall from last time

- Metric: *A description of the distance between two points*
- Metric in 3 spatial dimensions:

$$ds^2 = \frac{dx^2}{1 - \kappa x^2 / R^2} + x^2 d\Omega^2$$

Flat: $\kappa = 0, x = r$

Negative: $\kappa = -1, x = R \sinh(r / R)$

Positive: $\kappa = 1, x = R \sin(r / R)$

$$d\Omega^2 = d\theta^2 + \sin^2 \theta d\phi^2$$

Distance
(squared)

Curvature
radius

Metric in 4D space-time

- Minkowski metric (used in special relativity) :

$$ds^2 = -c^2 dt^2 + dr^2 + r^2 d\Omega^2$$

Time

Space

Distance in 4D space-time, without
curvature.

Geodesics

- Geodesic: The shortest path between two points
- In special/general relativity: the shortest path between two points in 4D space-time
- Photons follow the *null geodesic*, for which $ds^2=0$.

$$ds^2 = 0 \quad \Rightarrow$$


$$0 = -c^2 dt^2 + dr^2 + r^2 d\Omega^2 \quad \Rightarrow$$

$$c^2 dt^2 = dr^2 \quad (\text{for radial path}) \quad \Rightarrow$$

$$\frac{dr}{dt} = \pm c \quad (\text{i.e. photons move with speed } c)$$

Robertson-Walker metric

(a.k.a. Friedmann-Lemaître-Robertson-Walker metric, Friedmann-Robertson-Walker or Friedmann-Lemaître metric)



Friedmann



Lemaître



Robertson



Walker

The R-W metric is an exact solution to Einstein's field equation of general relativity, assuming isotropy and homogeneity (the cosmological principle)

It is extremely important for modern cosmology!

Robertson-Walker metric II

- Valid for Universe described by general relativity (4D space-time with curvature and expanding space)
- Note: Assumes cosmological principle to hold

$$ds^2 = -c^2 dt^2 + a(t)^2 \underbrace{\left[\frac{dx^2}{1 - \kappa x^2 / R^2} + x^2 d\Omega^2 \right]}$$

Same as in 3D metric with curvature

Flat: $\kappa = 0, x = r$

Negative: $\kappa = -1, x = R \sinh(r / R)$

Positive: $\kappa = 1, x = R \sin(r / R)$

$$d\Omega^2 = d\theta^2 + \sin^2 \theta d\phi^2$$

Coordinates in R-W metric

- t = proper time ('cosmic time')
- (x, θ, ϕ) or (r, θ, ϕ) = comoving coordinates

Remember :


Negative curvature ($\kappa = -1$) : $x = R \sinh(r / R)$

No curvature ($\kappa = 0$) : $x = r$


Positive curvature ($\kappa = 1$) : $x = R \sin(r / R)$

Comoving coordinates

t_1



t_2



In comoving coordinates:


$$d_{AB}(t_1) = d_{AB}(t_2)$$

Comoving distance


- Comoving distance: A distance that stays constant despite cosmic expansion, i.e. follows the Hubble flow
- Other similar expressions: Comoving volume (often a comoving cube), comoving observers, comoving density

Intermission: What is the *comoving* distance between A and B at $z=9$?

13.3 Gyr ago (redshift $z=9$)



Now (redshift $z=0$)




100 Mpc

Is the Universe Infinite?

- Geometry -

Consider a 2D world located in the surface layer of a 3D sphere




This 2D world will have finite area but no wall-like boundary

Is the Universe Infinite?

- Geometry -

Now, consider 3 spatial dimensions with curvature



Positive curvature ($\kappa=1$) \rightarrow
'closed geometry',
finite Universe

Negative curvature ($\kappa=-1$) \rightarrow
'open geometry',
infinite Universe

Zero curvature ($\kappa=0$) \rightarrow
'flat geometry',
infinite Universe

Is the Universe Infinite?

- Topology -

- Complications:

- Despite what is stated in many textbooks, the finiteness of space depends on *geometry* and *topology*.
- A flat or negatively curved Universe can be finite, if the topology is non-trivial.

Asteroids:


Classic arcade game by Atari, from 1979

<http://www.freeasteroids.org/>

Is the Universe Infinite?


- Topology -

Topology of the Universe: Describes how various parts of space are connected




Asteroids (1979): A simple of a non-trivial topology, giving a finite 2D Universe without any visible boundary (wall)

Suggestion for Literature Exercise: Topology of the Universe



Luminet's dodecahedron



Escher (1960)

Proper distance


Proper distance from observer ($r = 0$) to r :

$$d_p(t) = a(t) \int_0^r dr = a(t)r$$


- Proper distance between points A & B at time t : defined by spatial geodesic between A & B when scale factor is fixed at $a(t)$.
- Grows over time due to cosmic expansion, i.e. *not* a comoving distance

Intermission: What is the *proper* distance between A and B at $z=9$?

13.3 Gyr ago (redshift $z=9$)



Now (redshift $z=0$)



100 Mpc

Recession velocities I


- Rate of change of the proper distance:

$$\dot{d}_p = \dot{a}r = \frac{\dot{a}}{a} d_p$$
$$v_p(t_0) = H_0 d_p(t_0)$$

- This implies:

$$d_p > \frac{c}{H_0} \rightarrow$$
$$v_p(t_0) > c$$

Recession velocities II




- $v > c$ (!) at large distances (high z)
- But note: Not objects moving through space, but recession velocity due to expansion of space itself

Cosmic dynamics:

Four important equations

- Friedmann equation
- Fluid equation
- Acceleration equation
- Equation of state



Learning how to use these is one of the major goals of this course!

Cosmic dynamics:

Four important equations

What do you need them for?

- Calculating:
 $\rho(t), a(t), a'(t), a''(t), T(t), t(z), D(z)$
- Examples of applications:
 - Predicting fate of the Universe
 - Testing ages of astronomical objects (stars/galaxies) against cosmological models

The Friedmann equation

$$H \equiv \frac{\dot{a}}{a} \Rightarrow$$

$$H(t)^2 = \frac{8\pi G}{3c^2} \varepsilon(t) - \frac{\kappa c^2}{R_0^2} \frac{1}{a(t)^2}$$

At the current time :

$$H_0^2 = \frac{8\pi G}{3c^2} \varepsilon_0 - \frac{\kappa c^2}{R_0^2}$$

Critical density

Critical density \equiv The energy density required to make the Universe spatially flat

$$\kappa = 0 \quad \Rightarrow$$

$$\varepsilon_c(t) = \frac{3c^2}{8\pi G} H(t)^2$$

and

$$\varepsilon_{c,0} = \frac{3c^2}{8\pi G} H_0^2$$

Numerically, the critical energy density at the current time corresponds to 1 hydrogen atom per 200 liters, or 140 Msolar per kpc³

The Density Parameter

Dimensionless density parameter

$$\Omega(t) \equiv \frac{\varepsilon(t)}{\varepsilon_c(t)}$$

Rewriting the Friedmann equation:

$$1 - \Omega(t) = -\frac{\kappa c^2}{R_0^2 a(t)^2 H(t)^2}$$

or, at the current time :


$$1 - \Omega_0 = -\frac{\kappa c^2}{R_0^2 H_0^2}$$

Once more: Energy determines geometry (curvature)

Ω and Geometry

Energy
density

Curvature
(geometry)



$$\Omega_o > 1$$

$\kappa = +1$
Closed

$$\Omega_o < 1$$

$\kappa = -1$
Open

$$\Omega_o = 1$$

$\kappa = 0$
Flat


Why only 3 possible values for κ ?

The curvature κ always appears together with R_0 . You can always rescale R_0 to get $\kappa=0, +1$ or -1

$$1 - \Omega_0 = - \frac{\kappa c^2}{R_0^2 H_0^2}$$

Intermission: Where to measure?

If you wanted to assess the total density in some volume of the Universe, where would you measure this? How would you pick this volume?



The Fluid Equation

The Friedmann equation does not, by itself, allow a prediction of how the scale factor evolves with time. You also need the *fluid equation*.
Assumption: The energy components of the Universe can be treated as perfect fluids on large scales.

First law of Thermodynamics:

$$dQ = dE + PdV$$

Heat flow
into or out of
region

Change
in internal
energy of region

Change
in volume
of region

The Fluid Equation

Cosmic expansion is an adiabatic process,
giving no increase in entropy $\rightarrow dQ = 0$

Hence,

$$dE + PdV = 0$$

From this, you can derive the fluid equation:

$$\dot{\varepsilon} + 3\frac{\dot{a}}{a}(\varepsilon + P) = 0$$

The Acceleration Equation

The Friedmann equation and the fluid equation can be combined to form the acceleration equation:

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3c^2}(\varepsilon + 3P)$$

Important implication: Positive energy and pressure makes cosmic expansion slow down.
But a component with negative pressure ('tension') could cause the expansion to speed up.

The Equation of State

The equation of state relates the pressure and energy of the cosmic fluids:

$$P = w\varepsilon$$

Equations of states for the some of the most common components considered in cosmology:


$w = 0$ (non – relativistic matter)

$w = 1/3$ (radiation and relativistic matter)

$w = -1$ (cosmological constant)

$w < -1/3$ (dark energy) → Positive acceleration

The Cosmological Constant I



In 1917 Einstein introduced a constant Λ in his equations to produce a static matter-filled Universe.

After Hubble's discovery of cosmic expansion in 1929, Einstein referred to the introduction of Λ as his "greatest blunder"

The Cosmological Constant II

Introduction of $\Lambda \rightarrow$

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3c^2} \varepsilon(t) - \frac{\kappa c^2}{R_0^2} \frac{1}{a(t)^2} + \frac{\Lambda}{3}$$

$$\dot{\varepsilon} + 3\frac{\dot{a}}{a}(\varepsilon + P) = 0 \quad (\text{no change})$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3c^2}(\varepsilon + 3P) + \frac{\Lambda}{3}$$

The Cosmological Constant III

Λ corresponds to an energy of:

$$\varepsilon_{\Lambda} = \frac{c^2}{8\pi G} \Lambda$$

While Λ can in principle produce a static Universe ($a'=0$), this solution is as unstable as a pencil balanced on its sharpened point...

But: Because of its negative pressure, a Λ -like component with an energy slightly different than that assumed by Einstein has recently become part of the currently favoured cosmology, since this may explain the observed acceleration ($a''>0$) of the Universe