

Outline I

- What is dark matter?
- How much dark matter is there?
- How do we know it exists?
- Dark matter candidates
- The Cold Dark Matter (CDM) model

Outline II

- Dark halos and subhalos
- Problems with CDM
- Dark matter annihilation





Fritz Zwicky (1933): Dark matter in the Coma Cluster





How do we know that it exists?

- •Dynamics of galaxies
- •Dynamics and gas properties of galaxy clusters
- •Gravitational Lensing
- •Cosmological Parameters + Inventory of luminous material









Hot Gas in Galaxy Clusters

High mass required to keep the hot gas from leaving the cluster!

If gas in hydrostatic equilibrium \rightarrow Luminosity and temperature profile \rightarrow mass profile

X-ray gas, T=10⁷—10⁸ K





Intermission: One of these is not a lensed system - which one?





What is supersymmetry (SUSY)?

• A high-energy extension of the standard model

• SUSY predicts a symmetry between bosons and fermions:

Standard particle \leftrightarrow SUSY partner

fermion (e.g. quark) \leftrightarrow boson (e.g. squark)

boson (e.g. photon) \leftrightarrow fermion (e.g. photino) \rightarrow Zoo of new particles: selektrons, sneutrinos, gluinos,

Higgsinos, gravitinos, axinos...



Weakly Interacting Massive Particles (WIMPs)

- \bullet Interactions through weak force and gravity only \rightarrow dark matter transparent
- Weak-scale interactions \rightarrow right cosmological density to be dark matter ("The WIMP miracle")
- Massive (GeV to TeV scale)
- No WIMP candidate in standard model of particle physics
- The canonical WIMP is a SUSY particle (often a neutralino), but not all WIMP candidates are SUSYs





Generic assumptions (~100 GeV WIMPs) \rightarrow Handful of WIMPs in an average-sized coffee cup

Hot and Cold Dark Matter

- •Hot Dark Matter (HDM)
 - Relativistic early on (at decoupling)Ruled out by observations
- •Cold Dark Matter (CDM)
- •Non-relativistic early on (at decoupling) •The standard model for the non-baryonic
- dark matter •Successful in explaining the formation of large scale structure (galaxies, galaxy clusters, voids and filaments)

Additional Assumed CDM Properties

- Collisionless interacts *mainly* through gravity
- Dissipationless cannot cool by radiating photons
- Long-lived particles
- Behaves as perfect fluid on large scales











- Halo mass range: ~10⁻⁶ 10¹⁵ Msolar
 - Lower cutoff depends on detailed properties of the dark matter particles, could be 10⁻¹² to 10⁷ Msolar, depending on the model
 - •Mass function shape: Always far more low-mass halos than high-mass ones
 - •Low-mass halos assemble first, then merge to form high-mass ones





Subhalos

- Massive halos are assembled by the accretion of halos of lower mass
- Many accreted halos get disrupted in the tidal field of the halo they fell into, but some temporarily survive in the form of subhalos
- On average ~10% of the mass of a halo is in the form of subhalos at the current time









Density profiles of real galaxies I

• Singular Isothermal sphere

$$\rho_{\rm SIS}(r) = \frac{\rho(r_0)}{(r/r_0)^2}$$

$$\begin{split} \sigma(r) &= \text{constant} \\ \rho(r) &\to \infty \text{ when } r \to o \\ M(<r) &\to \infty \text{ when } r \to \infty \\ \text{Outer truncation required!} \end{split}$$

Works reasonably well for massive galaxies acting as strong gravitational lenses, probably due to baryon-domination in the centre





CDM problem II: Missing satellites

Possible solutions:

- Vanilla CDM incorrect alternative models (e.g. warm dark matter) produce fewer subhalos
- Star formation in low-mass subhalos inefficient → lots of ultrafaint or completely dark subhalos awaiting detection around the Milky Way

















Tidal dwarf galaxies

- TDGs form out of shredded disk material
- Only type of galaxy predicted and observationally confirmed to be nearly CDM-free







mechanism for creating galaxies without dark matter!

Nice topic for literature exercise!

Van Dokkum et al. 2018, Nature 555, 629